


Chemistry and target identification of natural products



22-24 May 2012, BUCHAREST, Romania

Aula of The Romanian Academy

Organizing Committee

CHAIR

Vasile I. PARVULESCU (RO)

INTERNATIONAL ADVISORY BOARD

Martin E. MAIER (DE)

Ari KOSKINEN (FI)

Roderich SÜSSMUTH (DE)

Stephen CLARK (UK)

Antti POSO (IT)

Camelia BALA (RO)

SECRETARIAT

Mihaela FLOREA (RO)

Bogdan COJOCARU (RO)

<http://www.unibuc.ro/COSTActionCM0804/>

Genome mining indicates that the genus *Xanthomonas* is a promising reservoir for new bioactive non-ribosomally synthesized peptides

Monique Royer¹, Ralf Koebnik², Mélanie Marguerettaz¹, Valérie Barbe³, Guillaume P. Robin², Chrystelle Brin⁴, Sébastien Carrere⁵, Camila Gomez¹, Manuela Hügelland⁶, Ginka Völler⁶, Julie Noël¹, Isabelle Pieretti¹, Saskia Rausch⁶, Valérie Verdier², Stéphane Poussier⁷, Philippe Rott¹, Roderich D. Süssmuth⁶, Stéphane Cociancich¹

¹CIRAD, UMR BGPI, F-34398 Montpellier Cedex5, France

²IRD, UMR RPB, F-34394 Montpellier Cedex, France

³Genoscope, Centre National de Séquençage, F-91057 Evry Cedex, France

⁴INRA, UMR IRHS, F-49071 Beaucouzé, France

⁵INRA, UMR LIPM, F-31326 Castanet-Tolosan Cedex, France

⁶Institut für Chemie, Technische Universität Berlin, D-10623 Berlin, Germany

⁷Université de la Réunion, UMR PVBMT, F-97715 Saint-Denis, La Réunion, France

stephane.cociancich@cirad.fr

Introduction

Xanthomonas is a large genus of Gram-negative bacteria that cause disease in hundreds of plant species. To date, the only known small molecule synthesized by non-ribosomal peptide synthesis (NRPS) in this genus is albicidin produced by *Xanthomonas albilineans* [1]. The DNA gyrase inhibitor albicidin is not only an important virulence factor but also a possible lead structure for novel antibiotics [2]. This study aims to estimate the biosynthetic potential of *Xanthomonas* spp. by *in silico* analyses of NRPS genes with unknown function recently identified in the sequenced genomes of *X. albilineans* and related species of *Xanthomonas*.

Experimental

In silico analyses were performed on NRPS genes present in the published finished genome sequence of *X. albilineans* strain GPE PC73 (accession n°: NC_013722.1), in the published draft genome sequence of *Xanthomonas oryzae* pv. *oryzae* strain X11-5A (accession n°: AFHK00000000.1), in the unpublished sequence of the 82-kb length region containing META-B in *X. oryzae* pv. *oryzae* strain BAI-3 and in the unpublished draft genome sequence of strain *Xanthomonas* spp. XaS3. Specificity of adenylation domains in NRPS and signatures were predicted using the web server NRPSpredictor2 [3].

Results and discussion

This study revealed four strains of the genus *Xanthomonas* possessing a novel homologous NRPS gene cluster, hereafter referred to as META-B. The sequence alignment of META-B gene clusters from *Xanthomonas* strains is indicative of the biosynthesis of lipopeptides or peptides linked to an other non-amino acid substrate, and involving the biosynthesis and incorporation of the non-proteinogenic amino acids Dpg (3,5-dihydroxyphenyl-glycine), Dab (2,4-diamino butyric acid), β -Hty (β -hydroxy-tyrosine) as well as of amino acid(s) of unknown identity (Figure 1; [4]). This study revealed that each sequence of the peptide is strain-specific. If the biosynthetic pathways are functional, these peptides may exert different biological functions in plant-bacteria interactions.

Conclusions

This extensive *in silico* study shows that the genus *Xanthomonas* constitutes a promising reservoir for new non-ribosomally synthesized peptides. Experimental elucidation of this

